



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore \frac{2}{u} \frac{\partial u}{\partial x} = -\frac{1}{s-x} + \frac{1}{x+y-s}. \quad \therefore \frac{\partial u}{\partial x} = \frac{u}{2} \frac{2s-2x-y}{(s-x)(x+y-s)} \dots (2),$$

$$\text{and } \frac{\partial u}{\partial y} = \frac{u}{2} \frac{2s-2y-x}{(s-y)(x+y-s)} \dots (3).$$

Equating (2) and (3) to 0, we have  $2s-2x-y=0$ ,  $2s-2y-x=0$ , whence  $x=\frac{2}{3}s$ ,  $y=\frac{2}{3}s$ ,  $z=2s-x-y=\frac{2}{3}s$ .

Hence the triangle is equilateral; and there is evidently a maximum.

309. Proposed by S. G. BARTON, Ph. D., Clarkson School of Technology.

In practical problems involving maxima and minima, it is really the greatest and least values of the function which are desired. Show why we can assume that the maximum is the greatest value and the minimum the least value under the conditions.

No solution of this problem has been received.

310. Proposed by C. N. SCHMALL, New York City.

Evaluate  $\int_0^\pi \frac{dx}{1-2a\cos x+a^2}$ . Edwards' *Integral Calculus for Beginners*, page 131, ex. 9, (iii). The answer given is  $\frac{\pi}{1-a^2}$ . Is this a complete answer to the question?

I. Solution by H. PRIME, Boston, Mass.

$$\begin{aligned} \int_0^\pi \frac{dx}{1+a^2-2a\cos x} &= \left[ \frac{2}{\pm(1-a^2)} \tan^{-1} \left( \pm \frac{1+a}{1-a} \tan \frac{x}{2} \right) \right]_0^\pi \\ &= \frac{(2n+1)\pi}{\pm(1-a^2)} - \frac{2n\pi}{\pm(1-a^2)} = \frac{\pi}{\pm(1-a^2)}. \end{aligned}$$

It should be noted that if  $a=\pm 1$ , the function  $1/(1+a^2-2a\cos x)$  becomes infinite for  $x=0$  or  $x=\pi$ . Hence, in this case, the integral has no meaning for the required limits, as they do not exist. The result also shows this for  $\pi \div (1-a^2) = \infty$ , when  $a=\pm 1$ .

Also solved by J. Scheffer, Francis Rust, A. M. Harding, and V. M. Spunar.

311. Proposed by WILMER THOMPSON, Senior, Drury College.

Solve the differential equation,

$$\left(\frac{dy}{dx}\right)^3 + x^3 = ax \left(\frac{dy}{dx}\right).$$

[From Forsythe's *Differential Equations*, p. 47.]